

Effect of Concussion History on Neurocognition and Neuromuscular Function of College Football Players

Altovise C. Scott, MS, ATC; Allison D. Varnell, MS, ATC; Gary B. Wilkerson, EdD, ATC; Shellie N. Acocello, PhD, ATC, Marisa A. Colston, PhD, ATC




BACKGROUND AND PURPOSE

- Sport-related concussions are a growing concern, the vast majority of which result from participation in football¹
- Post-concussion recovery of cognitive function, postural balance, and reaction time slower with concussion history²
- Recent evidence suggests long-term impairments, and risk for concussion recurrence and musculoskeletal injury^{3,4}
- Improvement in speed of peripheral motor responses to visual stimuli associated with decreased concussion rate⁵
- The purpose of our research was to identify any associations between concussion history, neurocognitive function, neuromuscular control, and core or lower extremity (Core/LE) sprain or strain among college football players

PARTICIPANT CHARACTERISTICS AND PROCEDURES

- 34 NCAA Division I-FCS football players: 17 cases with concussion history and 17 matched controls
 - Cases: Age 20.2 ± 1.3 yrs, Mass 100.72 ± 22.72 kg, Height 183.03 ± 7.05 cm
 - Controls: Age 19.4 ± 1.1 yrs, Mass 101.92 ± 19.31 kg, Height 185.27 ± 5.54 cm
- Neurocognitive composite scores obtained from computerized test (ImPACT, Pittsburgh, PA)
 - Verbal Memory: attention processes, learning, and memory
 - Visual Memory: visual attention and scanning, learning, and memory
 - Visual Motor Speed: visual processing, learning and memory, and visual-motor response speed
 - Composite Reaction Time (CRT): response speed (ms)
 - Impulse Control: measure of response errors
- Visual detection time (VDT) quantified by visuomotor testing system (Dynavision International, West Chester, OH)
 - Elapsed time (ms) between target illumination and initial motor response (release of depressed button)
 - Central and peripheral VDT assessed for right and left tests (Figure 1 A-D)
- Unilateral Squat Hold (USH): knee maintained in 45° flexion with heel elevated ~ 2.5 cm for 60 s (Figure 2)
 - Time to loss of postural balance recorded; balance quickly re-established for completion of test
 - USH postural balance also quantified by electronic tablet accelerometer application (Sway Medical, Tulsa, OK)
 - 15 cases and 15 controls (test data corrupted in 2 cases; corresponding matched control data excluded)
- Occurrence of Core/LE sprain or strain documented throughout pre-season practice sessions and 14-game season
 - Injury defined as any acute Core/LE sprain or strain that required evaluation and received treatment
- Receiver operating characteristic (ROC) analyses used to identify cut-points for dichotomization of variables
- Cross-tabulation analyses performed to assess associations between concussion history and dichotomous variables
- Logistical Regression analyses identified models that provided maximum discrimination between cases and controls

Figure 1



RESULTS

- ROC and cross-tabulation analyses identified 7 variables strongly associated with concussion history (Table 1)
 - Fisher's exact 1-sided *P*-value reported for categorizations created by cut-points derived from ROC analyses
- Multivariable 4-factor model:
 - USH (OR_{Adj} = 5.68), Peripheral VDT (OR_{Adj} = 3.23), Visual Memory (OR_{Adj} = 3.11), and CRT (OR_{Adj} = 2.27)
 - Model $\chi^2(4) = 11.04$; *P* = .026; Nagelkerke R² = .370
- Multivariable 3-factor model:
 - USH (OR_{Adj} = 7.31), Peripheral VDT (OR_{Adj} = 3.57), and Visual Memory (OR_{Adj} = 3.41)
 - Model $\chi^2(3) = 10.59$; *P* = .014; Nagelkerke R² = .357
- ROC analysis provided comparison of simplified 4-factor and 3-factor models (Figure 2)
 - Model comparisons: 4-factor ≥ 3 positive, 4-factor ≥ 2 positive, and 3-factor ≥ 2 positive (Table 2)
- Concussion history was strongly associated with the occurrence of Core/LE sprain or strain
 - Fisher's exact 1-sided *P* = .007; Sensitivity 79%; Specificity 70%; OR = 8.56 (90%CI; 2.24 – 32.63)

Table 1

Variable	Cut-Point	Sensitivity	Specificity	Odds Ratio	90% CI	P-Value
Visual Memory Score (0-100)	≤ 73.5	65%	71%	4.40	1.31 - 14.75	.042
Composite Reaction Time (ms)	≥ 675	29%	94%	6.67	0.99 - 44.94	.087
Impulse Control Score (0-100)	≥ 8	53%	65%	2.06	0.65 - 6.55	.245
Central VDT (ms)	≥ 270	82%	41%	3.27	0.87 - 12.27	.129
Peripheral VDT (ms)	≥ 298	82%	41%	3.27	0.87 - 12.27	.129
Postural Sway (Std Dev of m/s ²)	≥ .042	40%	87%	4.33	0.95 - 19.83	.107
Unilateral Squat Hold (s)	≤ 23	47%	88%	6.67	1.53 - 29.10	.029

Figure 2



Table 2


Prediction Model	Sensitivity	Specificity	Odds Ratio	90% CI	P-Value
3-Factor ≥2 Positive	77%	71%	7.80	2.16 – 28.19	.007
4-Factor ≥2 Positive	77%	71%	7.80	2.16 – 28.19	.007
4-Factor ≥3 Positive	47%	94%	14.22	2.18 – 92.69	.008

Table 3

Concussion History	Injury	No Injury
Yes	14	3
No	6	11
Total	20	14

OR = 8.56

Figure 3



CLINICAL RELEVANCE

- The results of our retrospective case-control analysis suggest that concussion may have long-lasting effects
 - Alternatively, pre-existing deficiencies in neural function may increase susceptibility to concussion occurrence
- The results of our prospective case-control analysis suggest that concussion history is a Core/LE injury risk factor
 - Whether a cause or an effect of concussion, deficiencies in neural function appear to be important to assess
 - Neurocognitive processes relating to detection, interpretation, and generation of responses to visual stimuli
 - Neuromuscular processes relating to maintenance of postural stability during single-leg support
- The potential for injury risk reduction through administration of screening tests and implementation of training activities for development of specific neurocognitive and neuromuscular adaptations needs thorough investigation

REFERENCES

1. McCrea M, et al. Acute effects and recovery time following concussion in collegiate football players. *JAMA*. 2003;290:2556-2563.
2. Covassin T, Moran R, Wilhelm K. Concussion symptoms and neurocognitive performance of high school and college athletes who incur multiple concussions. *Am J Sports Med*. 2013;41:2885-2889.
3. Nordström A, Nordström P, Ekstrand J. Sports-related concussion increases the risk of subsequent injury by about 50% in elite male football players. *Br J Sports Med*. 2014;48:1447-1450.
4. Powers KC, Kalmar JM, Cinnelli, ME. Recovery of static stability following a concussion. *Gait Posture*. 2014;39:611-614.
5. Clark JF, Graman P, Ellis JK, et al. An exploratory study of the potential effects of vision training on concussion incidence in football. *Optom Vis Perform* 2015;3:7-13.